



# MDwise

*A McLaren Company*

## PTSD and Panic Attacks

### Table of Contents

- Topic Overview
- Credits

### Topic Overview

---

Feeling afraid is a common symptom of post-traumatic stress disorder (PTSD).

Having intense fear that comes on suddenly could mean you're having a panic attack. This sudden fear may come without warning or without any obvious reason. Or a panic attack may happen when something reminds you of your trauma.

During a panic attack, you may be afraid of dying or afraid of losing control of yourself. It may seem like things happening around you aren't real. An attack usually lasts from 5 to 20 minutes but may last even longer, up to a few hours. You have the most anxiety about 10 minutes after the attack starts.

You may have physical symptoms, including:

- Chest pain.
- A fast or pounding heartbeat.
- Difficulty breathing.
- Dizziness, shaking, or trembling.
- Stomach pain or nausea.
- Sweating.
- Chills or hot flashes.
- Feeling like you're choking.

If you've had more than one panic attack, or if you feel worried about the next time a panic attack will happen, then you may have panic disorder. Worrying about future panic attacks can cause stress and interfere with your life. You may try to avoid things that bring back memories of your traumatic event.

### Getting treatment

Talk to your doctor or health professional if you've had panic attacks or if you think you may have panic disorder. You will work together to find the best way to treat the panic attacks and PTSD. Cognitive-behavioral therapy or medicine may help you have less fear. This can be used to treat both panic attacks and PTSD.

In cognitive-behavioral therapy, you learn relaxation techniques that can help you cope with the physical

symptoms of panic attacks. This therapy helps you understand how your thoughts and your reaction to your memories cause you to feel stress. You may do "exposure" exercises in which you focus on stressful memories until you can overcome your fearful reaction.

Antidepressants and other medicines also are used to treat panic disorder.

For more information, see the topics Post-Traumatic Stress Disorder and Panic Attacks and Panic Disorder.

## Credits

---

**Current as of: June 16, 2021**

**Author: Healthwise Staff**

**Medical Review: Adam Husney MD - Family Medicine**

**Kathleen Romito MD - Family Medicine**

**Jessica Hamblen PhD - Post Traumatic Stress Disorder**

---

Note: The "printer friendly" document will not contain all the information available in the online document. Some information (e.g. cross-references to other topics, definitions or medical illustrations) is only available in the online version.

---



This information does not replace the advice of a doctor. Healthwise, Incorporated, disclaims any warranty or liability for your use of this information.

© 1995-2022 Healthwise, Incorporated. Healthwise, Healthwise for every health decision, and the Healthwise logo are trademarks of Healthwise, Incorporated.